

Cellular Activation of Signaling ELISA:

CASE™ Kit for JNK T183/Y185

<u>Catalog Number</u>	<u>Contents</u>
FE-004	One CASE Kit for 96 cell-based assays

Description

The CASE™ Kit for JNK is designed to analyze JNK phosphorylation in cultured human and mouse cell lines. This cell-based ELISA kit quantifies the amount of activated (phosphorylated) JNK protein relative to total JNK protein. The JNK phosphorylation sites are threonine 183 and tyrosine 185. By determining JNK protein phosphorylation in your experimental model system, you can verify JNK pathway activation in your cell line and determine the effect of various treatments, inhibitors, or activators on the pathway.

More information on JNK protein from NCBI:

“The protein [JNK] is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which thought to be related to cytochrome c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation.”

Reference: <http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&val=20986493>

Materials Included

Please check the kit components immediately after you receive this package. SuperArray is not responsible for any missing items not reported within two (2) business days upon receipt.

Shelf Life: All reagents are stable at the recommended temperature for 6 months after receipt of the kit.

Enough reagents are provided for 96 assays.

Box 1: Assay and Detection Reagents (Store at 4°C)	Box 2: Primary Antibodies¹ (Store at -20°C)	
Components	Components	Dilution Factor
10X Antigen Retrieval Buffer ²	Anti-Phospho-Protein Specific Antibody	1:100
10X Washing Buffer ²	Anti-Pan-Protein Specific Antibody	1:100
Antibody Dilution Buffer		
Blocking Buffer, dry powder ³		
Cell Staining Solution		
Developing Solution		
Secondary Antibody ¹ (Dilute 1:16 before use.)		
Stop Solution		
Washing Buffer Reservoir (4)		

¹Each antibody stock must be freshly diluted by the appropriate factor into Antibody Dilution Buffer on the same day of the experiment. See the CASE™ Kit User Manual for more details.

²10X Washing Buffer and 10X Antigen Retrieval Buffer must be diluted to 1X with dH₂O before use.

³Just before use, dissolve powder in 30 ml 1X Washing Buffer by mixing well. Once dissolved, Blocking Buffer may be stored at 4 °C for up to two (2) weeks.

Shipping and Storage Conditions

Box 1 is shipped at either ambient or a refrigerated temperature and should be stored at 4°C.
Box 2 is shipped on dry ice or cold packs and should be stored at -20°C.

Related Products

Inhibitors for Functional Assays: Inhibitors of pathway activation that serve as a control for specificity.

For follow-up at the expression level:

RT² Profiler™ PCR Arrays and Oligo GEArray® Microarrays
(See www.superarray.com/ArrayList.php for complete listings)

RT² Real-Time™ PCR Primer Sets – available for every human, mouse, and rat gene
(See www.superarray.com/QRTsearch.php for our primer set search engine)

For knock down of gene expression:

SureSilencing™ shRNA Plasmids – available for every human, mouse, and rat gene
(See www.superarray.com/RNAisearch.php for complete listings)